

2017

# Region of Smooth Functions for Positive Solutions to an Elliptic Biological Model

Joon Hyuk Kang

*Andrews University*, kang@andrews.edu

Timothy Robertson

*Andrews University*

Follow this and additional works at: <https://digitalcommons.andrews.edu/pubs>



Part of the [Non-linear Dynamics Commons](#), and the [Numerical Analysis and Computation Commons](#)

---

## Recommended Citation

Hyuk Kang, Joon and Robertson, Timothy, "Region of Smooth Functions for Positive Solutions to an Elliptic Biological Model" (2017). *Faculty Publications*. 706.

<https://digitalcommons.andrews.edu/pubs/706>

This Article is brought to you for free and open access by Digital Commons @ Andrews University. It has been accepted for inclusion in Faculty Publications by an authorized administrator of Digital Commons @ Andrews University. For more information, please contact [repository@andrews.edu](mailto:repository@andrews.edu).

## REGION OF SMOOTH FUNCTIONS FOR POSITIVE SOLUTIONS TO AN ELLIPTIC BIOLOGICAL MODEL

Timothy Robertson<sup>1</sup>, Joon H. Kang<sup>2 §</sup>

<sup>1,2</sup>Department of Mathematics

Andrews University

Berrien Springs, MI. 49104, USA

---

**Abstract:** The non-existence and existence of the positive solution to the generalized elliptic model

$$\begin{aligned}\Delta u + g(u, v) &= 0 \text{ in } \Omega, \\ \Delta v + h(u, v) &= 0 \text{ in } \Omega, \\ u = v = 0 &\text{ on } \partial\Omega,\end{aligned}$$

were investigated.

**Key Words:** non-existence and existence of the solution, positive solution, generalized elliptic model

---

### 1. Introduction

The question in this paper concerns the existence of positive coexistence states when all growth rates are nonlinear and combined, more precisely, the existence of the positive steady state of

$$\begin{aligned}\Delta u + g(u, v) &= 0 \text{ in } \Omega, \\ \Delta v + h(u, v) &= 0 \text{ in } \Omega, \\ u = v = 0 &\text{ on } \partial\Omega,\end{aligned}$$

where  $\Omega$  is a bounded domain in  $R^N$  with smooth boundary  $\partial\Omega$ , and  $g, h \in C^2$  are such that  $g_{uu} < 0, h_{vv} < 0, g_{uv} > 0, h_{uv} > 0$ .

---

Received: March 29, 2017

Revised: September 16, 2017

Published: October 25, 2017

© 2017 Academic Publications, Ltd.

url: [www.acadpubl.eu](http://www.acadpubl.eu)

<sup>§</sup>Correspondence author

## 2. Preliminaries

In this section, we state some preliminary results which will be useful for our later arguments.

**Definition 2.1.** (upper and lower solutions)

$$\begin{cases} \Delta u + f(x, u) = 0 & \text{in } \Omega, \\ u|_{\partial\Omega} = 0 \end{cases} \quad (1)$$

where  $f \in C^\alpha(\bar{\Omega} \times R)$  and  $\Omega$  is a bounded domain in  $R^n$ .

(A) A function  $\bar{u} \in C^{2,\alpha}(\bar{\Omega})$  satisfying

$$\begin{cases} \Delta \bar{u} + f(x, \bar{u}) \leq 0 & \text{in } \Omega, \\ \bar{u}|_{\partial\Omega} \geq 0 \end{cases}$$

is called an upper solution to (1).

(B) A function  $\underline{u} \in C^{2,\alpha}(\bar{\Omega})$  satisfying

$$\begin{cases} \Delta \underline{u} + f(x, \underline{u}) \geq 0 & \text{in } \Omega, \\ \underline{u}|_{\partial\Omega} \leq 0 \end{cases}$$

is called a lower solution to (1).

**Lemma 2.1.** *Let  $f(x, \xi) \in C^\alpha(\bar{\Omega} \times R)$  and let  $\bar{u}, \underline{u} \in C^{2,\alpha}(\bar{\Omega})$  be respectively, upper and lower solutions to (1) which satisfy  $\underline{u}(x) \leq \bar{u}(x), x \in \bar{\Omega}$ . Then (1) has a solution  $u \in C^{2,\alpha}(\bar{\Omega})$  with  $\underline{u}(x) \leq u(x) \leq \bar{u}(x), x \in \bar{\Omega}$ .*

We also need some information on the solutions of the following logistic equations.

**Lemma 2.2.**

$$\begin{cases} \Delta u + uf(u) = 0 & \text{in } \Omega, \\ u|_{\partial\Omega} = 0, u > 0, \end{cases}$$

where  $f$  is a decreasing  $C^1$  function such that there exists  $c_0 > 0$  such that  $f(u) \leq 0$  for  $u \geq c_0$  and  $\Omega$  is a bounded domain in  $R^n$ .

If  $f(0) > \lambda_1$ , then the above equation has a unique positive solution. We denote this unique positive solution as  $\theta_f$ .

The main property about this positive solution is that  $\theta_f$  is increasing as  $f$  is increasing.

### 3. Existence and Nonexistence of Steady State

We consider

$$\begin{aligned} \Delta u + g(u, v) &= 0 \text{ in } \Omega \\ \Delta v + h(u, v) &= 0 \text{ in } \Omega \\ u = v = 0 &\text{ on } \partial\Omega, \end{aligned} \tag{2}$$

where  $\Omega$  is a bounded domain in  $R^N$  with smooth boundary  $\partial\Omega$  and  $g, h \in C^2$  are such that  $g_{uu} < 0, h_{vv} < 0, g_{uv} > 0, h_{uv} > 0, g(0, v) \geq 0, h(0, v) \geq 0$ .

We derive the following nonexistence result, which establishes a necessary condition for the existence of a positive solution to (2).

**Theorem 3.1.** *Suppose  $g_u(0, 0) > \lambda_1, h_v(0, 0) > \lambda_1$ , where  $\lambda_1$  is the first eigenvalue of  $-\Delta$  with homogeneous boundary condition, and there is  $c_0 > 0$  such that  $g_u(u, 0) < 0$  and  $h_v(0, v) < 0$  for  $u > c_0, v > c_0$ .*

(1) *If  $g_u(0, 0) \geq h_v(0, 0), -1 \leq g_{uu} < 0, h_{vv} \leq -1$  and*

$$\inf(h_{uv}) \inf(g_{uv}) + \inf(h_{uv}) + \inf(h_{vv}) \sup(h_{uv}) + \inf(h_{vv}) \geq 0,$$

*then (2) has no positive solution.*

(2) *If  $g_u(0, 0) \leq h_v(0, 0), -1 \leq h_{vv} < 0, g_{uu} \leq -1$  and*

$$\inf(g_{uv}) \inf(h_{uv}) + \inf(g_{uv}) + \inf(g_{uu}) \sup(g_{uv}) + \inf(g_{uu}) \geq 0,$$

*then (2) has no positive solution.*

*Proof.* Suppose the conditions in (1) or (2) holds and (2) has a positive solution  $(u, v)$ .

By the Mean Value Theorem, there is  $\bar{u}$  such that  $0 \leq \bar{u} \leq u$  and  $g(u, v) - g(0, v) = ug_u(\bar{u}, v)$ , and so by the monotonicity of  $g_u$

$$\begin{aligned} \Delta u + ug_u(u, v) &\leq \Delta u + ug_u(\bar{u}, v) \\ &= \Delta u + g(u, v) - g(0, v) \\ &= \Delta u + g(u, v) \\ &= 0. \end{aligned}$$

Similarly, we can prove that

$$\Delta v + vh_v(u, v) \leq 0.$$

Hence,  $(u, v)$  is an upper solution to

$$\begin{aligned} \Delta u + ug_u(u, v) &= 0 \text{ in } \Omega \\ \Delta v + vh_v(u, v) &= 0 \text{ in } \Omega \\ u = v = 0 &\text{ on } \partial\Omega. \end{aligned}$$

By the conditions  $(\tilde{u}, \tilde{v}) = (\theta_{g_u(\cdot, 0)}, \theta_{h_v(0, \cdot)})$  exist. We claim that for sufficiently small  $\epsilon > 0$ ,  $(\epsilon\tilde{u}, \epsilon\tilde{v})$  is a lower solution to

$$\begin{aligned} \Delta u + u g_u(u, v) &= 0 \text{ in } \Omega \\ \Delta v + v h_v(u, v) &= 0 \text{ in } \Omega \\ u = v = 0 &\text{ on } \partial\Omega. \end{aligned}$$

By the monotonicity of  $g_u$ , we have

$$\begin{aligned} \Delta(\epsilon\tilde{u}) + \epsilon\tilde{u}g_u(\epsilon\tilde{u}, \epsilon\tilde{v}) &\geq \Delta(\epsilon\tilde{u}) + \epsilon\tilde{u}g_u(\tilde{u}, 0) \\ &= \epsilon[\Delta(\tilde{u}) + \tilde{u}g_u(\tilde{u}, 0)] \\ &= 0. \end{aligned}$$

Similarly, we can prove that

$$\Delta(\epsilon\tilde{v}) + \epsilon\tilde{v}g_u(\epsilon\tilde{u}, \epsilon\tilde{v}) \geq 0.$$

Hence, we conclude that  $(\epsilon\tilde{u}, \epsilon\tilde{v})$  is a lower solution to

$$\begin{aligned} \Delta u + u g_u(u, v) &= 0 \text{ in } \Omega \\ \Delta v + v h_v(u, v) &= 0 \text{ in } \Omega \\ u = v = 0 &\text{ on } \partial\Omega. \end{aligned}$$

Therefore, by the Lemma 2.1, there is a positive solution to

$$\begin{aligned} \Delta u + u g_u(u, v) &= 0 \text{ in } \Omega \\ \Delta v + v h_v(u, v) &= 0 \text{ in } \Omega \\ u = v = 0 &\text{ on } \partial\Omega, \end{aligned}$$

which contradicts to the result in [1]. We now establish a sufficient condition for existence of a positive solution to (2).

**Theorem 3.2.** *Suppose  $g_u(0, 0) > \lambda_1, h_v(0, 0) > \lambda_1$ , and there are  $M > 0, N > 0$  such that  $g(M, N) < 0, h(M, N) < 0$ .*

*Then there is a positive solution to (2).*

*Proof.* By the condition, we have an upper solution  $(M, N)$  to (2). Let  $\phi$  be the first eigenfunction of  $-\Delta$  with homogeneous boundary condition. Then, by the continuity of  $g_u$  and  $h_v$  and the assumption that  $g_u(0, 0) > \lambda_1, h_v(0, 0) > \lambda_1$ ,  $g_u(\epsilon\phi, \epsilon\phi) > \lambda_1$  and  $h_v(\epsilon\phi, \epsilon\phi) > \lambda_1$  for sufficiently small  $\epsilon > 0$ .

By the Mean Value Theorem, there are  $\tilde{u}, \tilde{v}$  such that  $0 \leq \tilde{u} \leq \epsilon\phi, 0 \leq \tilde{v} \leq \epsilon\phi$  and

$$\begin{aligned} g(\epsilon\phi, \epsilon\phi) - g(0, \epsilon\phi) &= \epsilon\phi g_u(\tilde{u}, \epsilon\phi) \\ h(\epsilon\phi, \epsilon\phi) - h(\epsilon\phi, 0) &= \epsilon\phi h_v(\epsilon\phi, \tilde{v}). \end{aligned}$$

Hence, by the monotonicity of  $g_u$  and  $h_v$ ,

$$\begin{aligned} \Delta(\epsilon\phi) + g(\epsilon\phi, \epsilon\phi) &\geq \Delta(\epsilon\phi) + g(\epsilon\phi, \epsilon\phi) - g(0, \epsilon\phi) \\ &= \Delta(\epsilon\phi) + \epsilon\phi g_u(\tilde{u}, \epsilon\phi) \\ &\geq \epsilon(-\lambda_1\phi) + \epsilon\phi g_u(\epsilon\phi, \epsilon\phi) \\ &= \epsilon\phi[-\lambda_1 + g_u(\epsilon\phi, \epsilon\phi)] \\ &> 0, \end{aligned}$$

and

$$\begin{aligned} \Delta(\epsilon\phi) + h(\epsilon\phi, \epsilon\phi) &\geq \Delta(\epsilon\phi) + h(\epsilon\phi, \epsilon\phi) - h(\epsilon\phi, 0) \\ &= \Delta(\epsilon\phi) + \epsilon\phi h_v(\epsilon\phi, \tilde{v}) \\ &\geq \epsilon(-\lambda_1\phi) + \epsilon\phi h_v(\epsilon\phi, \epsilon\phi) \\ &= \epsilon\phi[-\lambda_1 + h_v(\epsilon\phi, \epsilon\phi)] \\ &> 0. \end{aligned}$$

Hence,  $(\epsilon\phi, \epsilon\phi)$  is a lower solution to (2). Therefore, by the Lemma 2.1, there is a positive solution to (2).

#### 4. Existence Region for Steady State

We consider

$$\begin{aligned} \Delta u + g(u, v) &= 0 \text{ in } \Omega \\ \Delta v + h(u, v) &= 0 \text{ in } \Omega \\ u = v = 0 &\text{ on } \partial\Omega, \end{aligned} \tag{3}$$

where  $\Omega$  is a bounded domain in  $R^N$  with smooth boundary  $\partial\Omega$  and  $g, h \in C^2$ .

We prove the following existence results.

**Theorem 4.1.** *Suppose  $g_u(0, 0) > \lambda_1, g(0, v) \geq 0, g_{uu} < 0, g_{uv} > 0$  and there is  $c_0 > 0$  such that  $g_u(u, 0) < 0, g(u, v) < 0$  for  $u > c_0, v > c_0$ . [ $h_v(0, 0) > \lambda_1, h(u, 0) \geq 0, h_{vv} < 0, h_{uv} > 0$  and there is  $c_0 > 0$  such that  $h_v(0, v) < 0, h(u, v) < 0$  for  $u > c_0, v > c_0$ .] Then there is a number  $M(g) < \lambda_1$  [ $N(h) < \lambda_1$ ] such that for any  $h \in C^2$  such that  $h(u, 0) \geq 0, h_{uv} > 0, h_{vv} < 0, h_v(0, v) < 0, h(u, v) < 0$  for  $u > c_0, v > c_0$  and  $h_v(0, 0) > M(g)$  [for any  $g \in C^2$  such that  $g_{uu} < 0, g_{uv} > 0, g_u(u, 0) < 0, g(u, v) < 0$  for  $u > c_0, v > c_0$ , and  $g_u(0, 0) > N(h)$ ], (3) has a positive solution  $u^+, v^+$  in  $\Omega$ .*

*Proof.* Let  $\underline{u} = \theta_{g_u(\cdot, 0)}$  be the unique positive solution to

$$\begin{aligned} \Delta u + u g_u(u, 0) &= 0 \text{ in } \Omega \\ u &= 0 \text{ on } \partial\Omega. \end{aligned}$$

Let  $M(g) = \lambda_1(-h_v(\theta_{g_u(\cdot,0)}, 0))$  be the smallest eigenvalue of

$$\begin{aligned} -\Delta Z - (h_v(\theta_{g_u(\cdot,0)}, 0) - h_v(0,0))Z &= \mu Z \text{ in } \Omega \\ Z &= 0 \text{ on } \partial\Omega. \end{aligned}$$

and  $\omega_0(x)$  be the corresponding normalized positive eigenfunction. By the monotonicity,  $M(g) < \lambda_1$ . Let  $\underline{v} = \epsilon\omega_0(x)$ . Let  $h \in C^2$  be such that  $h_{uv} > 0, h_{vv} < 0, h_v(0, v) < 0, h(u, v) < 0$  for  $u > c_0, v > c_0$  and  $h_v(0, 0) > M(g)$ . Then, by the Mean Value Theorem, there is  $\tilde{u}$  and  $\tilde{v}$  such that

$$\begin{aligned} 0 &\leq \tilde{u} \leq \underline{u} \\ 0 &\leq \tilde{v} \leq \underline{v} \\ g(\underline{u}, \underline{v}) - g(0, \underline{v}) &= \underline{u}g_u(\tilde{u}, \underline{v}) \\ h(\underline{u}, \underline{v}) - h(\underline{v}, 0) &= \underline{v}h_u(\underline{u}, \tilde{v}), \end{aligned}$$

so by the monotonicity of  $g_u$  and  $h_v$ , for sufficiently small  $\epsilon > 0$ ,

$$\begin{aligned} \Delta \underline{u} + g(\underline{u}, \underline{v}) &\geq \Delta \underline{u} + g(\underline{u}, \underline{v}) - g(0, \underline{v}) \\ &= \Delta \underline{u} + \underline{u}g_u(\tilde{u}, \underline{v}) \\ &\geq \Delta \underline{u} + \underline{u}g_u(\underline{u}, \underline{v}) \\ &= \Delta \underline{u} + \underline{u}[g_u(\underline{u}, 0) + g_u(\underline{u}, \underline{v}) - g_u(\underline{u}, 0)] \\ &= \underline{u}[g_u(\underline{u}, \underline{v}) - g_u(\underline{u}, 0)] \\ &> 0 \text{ in } \Omega \end{aligned}$$

and

$$\begin{aligned} \Delta \underline{v} + h(\underline{u}, \underline{v}) &\geq \Delta \underline{v} + h(\underline{u}, \underline{v}) - h(\underline{u}, 0) \\ &= \Delta \underline{v} + \underline{v}h_v(\underline{u}, \tilde{v}) \\ &\geq \Delta \underline{v} + \underline{v}h_v(\underline{u}, \underline{v}) \\ &= \Delta(\epsilon\omega_0) + \epsilon\omega_0 h_v(\theta_{g_u(\cdot,0)}, \epsilon\omega_0) \\ &= \Delta(\epsilon\omega_0) + \epsilon\omega_0 [h_v(\theta_{g_u(\cdot,0)}, 0) + h_v(\theta_{g_u(\cdot,0)}, \epsilon\omega_0) - h_v(\theta_{g_u(\cdot,0)}, 0)] \\ &= \epsilon [h_v(0, 0)\omega_0 - M(g)\omega_0] + \epsilon\omega_0 [h_v(\theta_{g_u(\cdot,0)}, \epsilon\omega_0) - h_v(\theta_{g_u(\cdot,0)}, 0)] \\ &\geq \epsilon\omega_0 [h_v(0, 0) - M(g)] + \epsilon^2\omega_0^2 \inf(h_{vv}) \\ &> 0 \text{ in } \Omega. \end{aligned}$$

So,  $\underline{u} > 0, \underline{v} > 0$  is a lower solution to (3). But, by the condition, there is a sufficiently large upper solution to (3). Therefore, there is a positive solution  $u^+, v^+$  of (3).

For the next Theorem, we set

$$S_g = \{h \in C^2 | h_{uv} > 0, M \leq h_{vv} < 0, h(u, 0) \geq 0,$$

there is  $c_0 > 0$  such that  $h(u, v) < 0$  for  $u > c_0, v > c_0\}$  for  $g \in C^2$  such that  $g_{uu} < 0, g_{uv} > 0, g(0, v) \geq 0$ , there is  $c_0 > 0$  such that  $g(u, v) < 0$  for  $u > c_0, v > c_0$  and

$$S_h = \{g \in C^2 | N \leq g_{uu} < 0, g_{uv} > 0, g(0, v) \geq 0,$$

there is  $c_0 > 0$  such that  $g(u, v) < 0$  for  $u > c_0, v > c_0\}$  for  $h \in C^2$  such that  $h_{uv} > 0, h_{vv} < 0, h(u, 0) \geq 0$ , there is  $c_0 > 0$  such that  $h(u, v) < 0$  for  $u > c_0, v > c_0$ .

**Theorem 4.2.** *Let  $g \in C^2$  such that  $g_{uu} < 0, g_{uv} > 0, g(0, v) \geq 0$ , there is  $c_0 > 0$  such that  $g(u, v) < 0$  for  $u > c_0, v > c_0$  and  $g_u(0, 0) \leq \lambda_1$  [ $h \in C^2$  such that  $h_{uv} > 0, h_{vv} < 0, h(u, 0) \geq 0$ , there is  $c_0 > 0$  such that  $h(u, v) < 0$  for  $u > c_0, v > c_0$  and  $h_v(0, 0) \leq \lambda_1$ ]. Then there is a number  $M(g) > \lambda_1$  [ $N(h) > \lambda_1$ ] such that for any  $h \in S_g$  satisfying  $h_v(0, 0) > M(g)$  [for any  $g \in S_h$  satisfying  $g_u(0, 0) > N(h)$ ], (3) has a positive solution in  $\Omega$ .*

*Proof.* Suppose  $g_u(0, 0) \leq \lambda_1$ . Let  $h \in S_g$  be such that  $h_v(0, 0) > \lambda_1$ . Since

$$\begin{aligned} \lim_{c \rightarrow \infty} \lambda_1(-g_u(0, \theta \frac{c}{-M}) + g_u(0, 0)) &\leq \lim_{c \rightarrow \infty} \lambda_1(-\inf(g_{uv})\theta \frac{c}{-M} + g_u(0, 0)) \\ &\leq \lim_{c \rightarrow \infty} \lambda_1(-\inf(g_{uv})\frac{c - \lambda_1}{-M}\phi_0 + g_u(0, 0)) \\ &= -\infty, \end{aligned}$$

there is a number  $M(g) \geq \lambda_1$  such that  $\lambda_1(-g_u(0, \theta \frac{c}{-M}) + g_u(0, 0)) < g_u(0, 0)$  if  $c > M(g)$ . Hence, if  $h_v(0, 0) > M(g)$ , then  $\lambda_1(-g_u(0, \theta \frac{h_v(0,0)}{-\inf(h_{vv})}) + g_u(0, 0)) < \lambda_1(-g_u(0, \theta \frac{h_v(0,0)}{-M}) + g_u(0, 0)) < g_u(0, 0)$ .

Let  $h_v(0, 0) > M(g)$  and  $\underline{u} = \epsilon\omega_0$  and  $\underline{v} = \theta \frac{h_v(0,0)}{-\inf(h_{vv})}$ , where  $\omega_0$  is the normalized positive eigenfunction corresponding to  $\lambda_1(-g_u(0, \theta \frac{h_v(0,0)}{-\inf(h_{vv})}) + g_u(0, 0))$ . Then by the Mean Value Theorem, there are  $\tilde{u}, \tilde{v}$  such that  $0 \leq \tilde{u} \leq \underline{u}, 0 \leq \tilde{v} \leq \underline{v}$  and

$$\begin{aligned} g(\underline{u}, \underline{v}) - g(0, \underline{v}) &= \underline{u}g_u(\tilde{u}, \underline{v}) \\ h(\underline{u}, \underline{v}) - h(\underline{u}, 0) &= \underline{v}h_v(\underline{u}, \tilde{v}). \end{aligned}$$



Hence, by the monotonicity of  $g_u$  and  $h_v$ , for sufficiently small  $\epsilon > 0$ ,

$$\begin{aligned}
 & \Delta \underline{u} + g(\underline{u}, \underline{v}) \\
 \geq & \Delta \underline{u} + g(\underline{u}, \underline{v}) - g(0, \underline{v}) \\
 = & \Delta \underline{u} + \underline{u}g_u(\tilde{u}, \underline{v}) \\
 \geq & \Delta \underline{u} + \underline{u}g_u(\underline{u}, \underline{v}) \\
 = & \Delta \underline{u} + \underline{u}[g_u(0, 0) + g_u(\underline{u}, \underline{v}) - g_u(0, \underline{v}) + g_u(0, \underline{v}) - g_u(0, 0)] \\
 \geq & \Delta \underline{u} + \underline{u}[g_u(0, 0) + \inf(g_{uu})\underline{u} + g_u(0, \underline{v}) - g_u(0, 0)] \\
 = & \Delta(\epsilon\omega_0) + \epsilon\omega_0[g_u(0, 0) + \inf(g_{uu})\epsilon\omega_0 + g_u(0, \theta_{\frac{h_v(0,0)}{-\inf(h_{vv})}}) - g_u(0, 0)] \\
 = & -\epsilon\lambda_1[-g_u(0, \theta_{\frac{h_v(0,0)}{-\inf(h_{vv})}}) + g_u(0, 0)]\omega_0 + g_u(0, 0)\epsilon\omega_0 + \epsilon^2\omega_0^2 \inf(g_{uu}) \\
 = & \epsilon\omega_0[g_u(0, 0) - \lambda_1(-g_u(0, \theta_{\frac{h_v(0,0)}{-\inf(h_{vv})}}) + g_u(0, 0))] + \epsilon^2\omega_0^2 \inf(g_{uu}) \\
 > & 0 \text{ in } \Omega
 \end{aligned}$$

and

$$\begin{aligned}
 & \Delta \underline{v} + h(\underline{u}, \underline{v}) \\
 \geq & \Delta \underline{v} + h(\underline{u}, \underline{v}) - h(\underline{u}, 0) \\
 = & \Delta \underline{v} + \underline{v}h_v(\underline{u}, \tilde{v}) \\
 \geq & \Delta \underline{v} + \underline{v}h_v(\underline{u}, \underline{v}) \\
 = & \Delta \underline{v} + \underline{v}[h_v(0, 0) + h_v(\underline{u}, \underline{v}) - h_v(\underline{u}, 0) + h_v(\underline{u}, 0) - h(0, 0)] \\
 \geq & \Delta \underline{v} + \underline{v}[h_v(0, 0) + \inf(h_{vv})\underline{v} + h_v(\underline{u}, 0) - h_v(0, 0)] \\
 = & \underline{v}[h_v(\underline{u}, 0) - h_v(0, 0)] \\
 > & 0 \text{ in } \Omega.
 \end{aligned}$$

So,  $\underline{u}, \underline{v}$  is a lower solution to (3). Hence, by the condition, if  $h_v(0, 0) > M(g)$ , there is a positive solution to (3).

### References

- [1] B. Chase, J. Kang, Positive solutions to an elliptic biological model, *Global Journal of Pure and Applied Mathematics*, **5**, No. 2 (2009), 101-108.
- [2] P. Korman, A. Leung, On the existence and uniqueness of positive steady states in the Volterra-Lotka ecological models with diffusion, *Appl. Anal.*, **26**, No. 2 (1987), 145-160.
- [3] L. Zhengyuan, P. De Mottoni, Bifurcation for some systems of cooperative and predator-prey type, *J. Partial Differential Equations* (1992), 25-36.